

# Quantum Interferometry and Gravitation with Positronium

Marco G. Giammarchi

*Istituto Nazionale Fisica Nucleare Sezione di Milano*



## QUPLAS

QUantum Interferometry, decoherence and gravitational studies with Positrons and LASers

- Outline of talk:**
- Theoretical motivation
  - Talbot Quantum Interferometry
  - Design and test of QUPLAS-0
  - QUPLAS-I and Ps Interferometry
  - Positronium fall (QUPLAS-II)

Home of the Experiment:  
L-NESS Laboratory of the  
Milano Politecnico in Como

# The QUPLAS Collaboration

## Università degli Studi di Milano and Infn Milano

S. Castelli, S. Cialdi, E. Dei Cas, M. Giammarchi\*, M. Longhi, G. Maero  
L. Miramonti, S. Olivares, M. Paris, M. Potenza, M. Romè, S. Sala



## Politecnico di Milano (at Como)

S. Aghion, M. Bollani (INFN del CNR), C. Evans, M. Leone, R. Ferragut



## Albert Einstein Center – Laboratory for HEP – Bern University

A. Ariga, T. Ariga, A. Ereditato, C. Pistillo, P. Scampori



## Dep.t of Chemistry, University of Bath

K. Edler



R. Greaves (Los Angeles, formerly at First Point Scientific)

# Ps : the truly elementary atom


## Energy levels of hydrogen and positronium

$$E_n(H) = -\frac{\mu e^4}{2\hbar n^2} = -\frac{1}{n^2} \times 13.6 \text{ eV}$$

$$\mu_H = \frac{m_e M}{m_e + M} \approx m_e$$

$$\mu_{Ps} = \frac{m_e^2}{2m_e} = \frac{m_e}{2}$$

$$E_n(Ps) = -\frac{1}{n^2} \times 6.8 \text{ eV}$$



A pure QED system where spin-orbit and hyperfine effects are of the same order

The metastable electron-positron bound state can exist in different configurations depending on the relative spin states of the positron and the electron. These are known as para-positronium (p-Ps), with total spin  $S = 0$  and ortho-positronium (o-Ps) with  $S = 1$ .

These spin states have very different lifetimes:

$$|S, m\rangle = |0, 0\rangle = \frac{1}{\sqrt{2}} (|\uparrow\downarrow\rangle - |\downarrow\uparrow\rangle)$$

$$\tau_{p\text{-Ps}} = 125 \text{ ps}$$

$$|S, m\rangle = |1, 0\rangle = \frac{1}{\sqrt{2}} (|\uparrow\downarrow\rangle + |\downarrow\uparrow\rangle)$$

$$\tau_{o\text{-Ps}} = 142 \text{ ns}$$

$$|S, m\rangle = |1, 1\rangle = |\uparrow\uparrow\rangle$$

$$|S, m\rangle = |1, -1\rangle = |\downarrow\downarrow\rangle$$

Any process that converts o-Ps to p-Ps is easy to see in lifetime spectra

## 1951: First production of positronium by Martin Deutsch





FIG. 1. Decay curves of positronium in several gases. The dotted lines are corrected for time resolution of the instrument.

100 ns lifetime of triplet Ps



From M. Deutsch  
Phys. Rev. **82**, 455 (1951)

### Separation of Ps from Radioactive source



## Our systems of interest :


- Electron (an elementary fermion)
- Positron (the antifermion)
- Positronium (Ps, a particle/antiparticle symmetric system)

# Introduction to the concept of Quantum Interferometry of Ps

## The typical structure of a Quantum Mechanical Experiment

### Preparation :

- $e^+$  beam
- Ps beam
- Target
- Laser (excitation)
- First grating



### Detection :

- Recording interference pattern
- Projection on measurement eigenstates

Preparation

Detection

Non – ideality :  
Incoherence

Interaction  
Propagation  
Interference

Non – ideality :  
Decoherence

# Theoretical motivation

Here is a (Lee Smolin's inspired) arbitrary list of outstanding problems in Theoretical, Particle Physics and Cosmology that are related to QUPLAS:

- **Theoretical motivation**
- Talbot Quantum Interferometry
- Design and test of QUPLAS-0
- QUPLAS-I and Ps Interferometry
- Positronium fall (QUPLAS-II)

**The problem of Quantum Gravity:** Combine General Relativity and Quantum Theory into a single theory that can claim to be the complete theory of nature.

**The foundational aspects of Quantum Mechanics:** Address the interpretational and epistemological aspects in the foundations of Quantum Mechanics.

**The unification of particles and forces:** Determine whether or not the various particles and forces can be unified in a theory that explains them all as manifestations of a single, fundamental entity.

**The problem of quantum gravity:** Combine General Relativity and Quantum Theory into a single theory that can claim to be the complete theory of nature.

**The foundational aspects of quantum mechanics:** Address the interpretational and epistemological aspects in the foundations of Quantum Mechanics.

**The unification of particles and forces:** Determine whether or not the various particles and forces can be unified in a theory that explains them all as manifestations of a single, fundamental entity.

QUPLAS-0) CPT test on fundamental fermions

QUPLAS-I) Test of Decoherence and of the Born Interpretation of the Wave Function

QUPLAS-II) Test of the Weak Equivalence Principle (WEP) for the simplest atom: positronium.

## Gravity and the Particles

**General Relativity** is the current Theory of Gravity

It is a classical (geometric) theory

It has never been tested at the level of elementary particles

It has no quantum version (approximations in weak field can be quantized with graviton)

From the particle physics point of view, it could be mediated by a tensor (spin-2) carrier

# Gravity and the Particles (CPT)

Dynamical meaning

$$F = m_I a$$

The gravitational «charge»

$$F = -G m_G M_G / r^2$$

According to the WEP

$$m_I = m_G$$

CPT Theorem


$$m_I = \bar{m}_I$$

$$m_G = m_I = \bar{m}_I ? \bar{m}_G$$

Which means that

$$m_G \neq \bar{m}_G$$


Would not mean that CPT is broken



# Gravity and the Particles

In many Quantum Gravity models (in the classical static limit), one has :

$$V = -GmM / r \left( 1 \mp a e^{-r/v} + b e^{-r/s} \right)$$



- The sign of the Gravi-Vector can be different between Matter and Antimatter
- Ranges and strength unknowns

From the Particle Physics point of view, it could be mediated by a tensor (spin-2) carrier, with the charge being mass-energy.

	<b>Matter-Matter (e- e-)</b>	<b>Antimatter-Matter (e+ e-)</b>	<b>Quantum Gravity</b>
Scalar	attractive	attractive	gravi-scalar
Vector	repulsive	attractive	gravi-vector
Tensor (Gravity)	attractive	attractive	graviton
Tensor (Antigravity)	attractive	Repulsive (CPT violating)	



# Gravity and the Particles

## Experimental tests of the Weak Equivalence Principle

Where do we stand ?

### Matter

- Weak Equivalence Principle tested on many different systems
- Torsion Balance Measurement
- $10^{-13}$  level reached

### Antimatter


- $g$  not measured
- Antihydrogen program at CERN (e. g. The AEGIS experiment)
- Aiming at 1% accuracy

### Positronium

- Matter/Antimatter system

?

# Talbot Quantum Interferometry



- Theoretical motivation
- **Talbot Quantum Interferometry**
- Design and test of QUPLAS-0
- QUPLAS-I and Ps Interferometry
- Positronium fall (QUPLAS-II)

Period  $D$     Slit width  $a$   
Open fraction  $a/D$

De Broglie wave impinging on a grating

- Classical propagation in the  $y$  direction
- Schroedinger dynamics in the  $x$  direction
- Neglect  $z$ -axis diffraction

Interference on the screen: «Fresnel» integral


$$\psi^{(N)}(x, t=0) \approx \sum_{n=1}^N \psi_n(x, t=0)$$

$$H_{\text{eff}} = \frac{p_x^2}{2m}$$

$$I(x) = \left| \psi^{(N)}(x, t = L/v) \right|^2$$

$$\psi^{(N)}(x, t) = \frac{1}{\sqrt{\lambda L}} \int_{-\infty}^{+\infty} \exp\left[ i \frac{\pi}{\lambda L} (x - x')^2 \right] \psi^{(N)}(x', 0) dx'$$

# Talbot and Fraunhofer




Considering the usual expansion, one can define the F parameter

$$F = k \frac{\bar{x}'^2}{\bar{x}^2} = \frac{2\pi}{\lambda} \frac{\bar{x}'^2}{\bar{x}^2} \approx \frac{1}{\lambda} \frac{a^2}{L}$$

$$|\bar{x} - \bar{x}'| = \sqrt{\bar{x}'^2 + \bar{x}^2 - 2\bar{x}\bar{x}'} = |\bar{x}| \sqrt{1 + \frac{\bar{x}'^2}{\bar{x}^2} - 2\frac{\bar{x}\bar{x}'}{\bar{x}^2}}$$

$F \ll 1$  in the Fraunhofer case  
 $F > 1$  in the Talbot case

# The Talbot-Lau Effect vs Fraunhofer



$$L_T = \frac{a^2}{\lambda}$$

Talbot length

The Talbot case


Generates coherence

Interference

G1 revival

$$L = L_T$$

Pattern at C



The Fraunhofer case

$$L \gg L_T$$

$$(BC \gg L_T)$$


# Talbot carpets

The characteristic pattern of the Talbot effect can be used to make sure the observed effect is the Talbot effect for the specified wavelength

Units  
Talbot  
length


Fringes visibility for the given  
wavelength

$$c = \frac{I_{\max} - I_{\min}}{I_{\max} + I_{\min}}$$



«Ballistic» moiré  
regime

Fraunhofer regime  
setting in when  $L \gg L_T$




# Design and test of QUPLAS-0

## The Como continuous e+ beam


The VEPAS Laboratory at the L-Ness Politecnico di Milano at Como (R. Ferragut).

<http://www.como.polimi.it/positron>

- Theoretical motivation
- Talbot Quantum Interferometry
- **Design and test of QUPLAS-0**
- QUPLAS-I and Ps Interferometry
- Positronium fall (QUPLAS-II)



**Slow positron beam.** 1. Radioactive source; 2. Electrostatic optics; 3. Sample chamber; 4. HpGe detectors; 5. Cryostat; 6. High voltage protection cage; 7. Power suppliers; 8. Detector electronics.



Original intensity of the source: 50 mCi (current : ~ 10 mCi)  
Tungsten moderator  $\rightarrow$  reduces E down to a few eV  
Electrostatic transport  $\rightarrow$  positron beam

# QUPLAS - 0


- Interferometry with positrons
- Interferometry with electrons (in the same apparatus)
- Comparison
- A new CPT test

The positron/electron beam :

$T = 10$  keV (typical)

The gratings ( $\sim \mu\text{m}$  thickness):

SiNx based substrates  
Electron Beam Litography



The detector :  
Nuclear Emulsions ( $\sim \mu\text{m}$  resolution) over a potentially large (mm or cm size) area.

Demonstrated for  $p$ -bar in AEGIS at CERN

S. Aghion et al., JINST 8 (2013) P08013.

Positron beam energy: from a few keV up to 20 keV

Reference value: 10 keV

Intensity:  $\sim 4 \times 10^4$  e<sup>+</sup>/s

$$T = 10 \text{ keV} \quad v = 6 \times 10^7 \text{ m/s}$$

The de Broglie wavelength

$$\lambda = \frac{h}{mv} = 1.2 \times 10^{-11} \text{ m}$$


Given a grating with

$$d = 2 \text{ } \mu\text{m}$$

The Talbot length

$$L_T = \frac{d^2}{\lambda} = 33 \text{ cm}$$

QUPLAS-0



One can choose  $b = c = 33$  cm


To have a  $2 \text{ } \mu\text{m}$  periodicity pattern on C

- Setup preparation
- Exposure to the e<sup>+</sup> beam
- Integration on the emulsion detector C





# QUPLAS – 0 : emulsion exposure to 9-18 keV e+ beam

New type of emulsions produced at LHEP




Scanning with the LHEP microscopes driven by ad hoc software (in Bern)




120 s integration of the QUPLAS-0 beam profile for 15 keV e+

Colors are relative to number of positrons per  $10^4 \mu\text{m}^2$  emulsion surface

The nearly gaussian positron spot of FWHM ( $\sim 2.4$  mm) was analyzed in the emulsion in an area of  $1.5 \times 1.5$  cm<sup>2</sup>



Number of counts observed as a function of energy, when normalized taking into account the effect of the (1 μm thick) protective layer shows the high intrinsic emulsion efficiency to e<sup>+</sup>'s at energies from 9 to 18 keV.




S. Aghion et al, J. of Instr. 11 (2016) P06017

# QUPLAS – 0 : building up a test interferometer

Interferometer frame

First grating holder



Second grating

Emulsion holder





# QUPLAS-I and Ps Interferometry

Problems to face :


- Positronium is a neutral atom
- Positronium has a very short lifetime

Detection through ionization required.


Laser excitation required.

- Theoretical motivation
- Talbot Quantum Interferometry
- Design of QUPLAS-0
- **QUPLAS-I and Ps Interferometry**
- Positronium fall (QUPLAS-II)

QUPLAS-I and  
QUPLAS-II Interferometry



# Positrons and Positronium (Ps)



ortho-Ps is short lived

But its lifetime can be increased by exciting it on a Rydberg (high-n) state

$$\tau \approx n^3 l^2$$

The metastable electron-positron bound state can exist in different configurations depending on the relative spin states of the positron and the electron. These are known as para-positronium (p-Ps), with total spin  $S = 0$  and ortho positronium (o-Ps) with  $S = 1$ .

These spin states have very different lifetimes:

$$|S, m\rangle = |0, 0\rangle = \frac{1}{\sqrt{2}} (|\uparrow\downarrow\rangle - |\downarrow\uparrow\rangle)$$

$$\tau_{p\text{-Ps}} = 125 \text{ ps}$$

$$|S, m\rangle = |1, 0\rangle = \frac{1}{\sqrt{2}} (|\uparrow\downarrow\rangle + |\downarrow\uparrow\rangle)$$

$$\tau_{o\text{-Ps}} = 142 \text{ ns}$$


$$|S, m\rangle = |1, 1\rangle = |\uparrow\uparrow\rangle$$

$$|S, m\rangle = |1, -1\rangle = |\downarrow\downarrow\rangle$$


A  
o  
ir

# Samples.


## Micrometre membranes



Synthesis route to the growth of free-standing surfactant-template films of silica (Chemistry Department, University of Bath, Claverton Down, Bath) [1]



Calcination ( $400^\circ\text{C}$ )



[1] K.J. Edler\* and B. Yang, *Chem. Soc. Rev.*, 2013 **42**, 3765

# QUPLAS – 0 and QUPLAS – I summary

## Positronium Quantum Interferometry concept

- Positron Interferometry
- Electron Interferometry
- Positronium Interferometry

An elementary fermion  
The relevant antifermion

The bound fermion-antifermion system  
(also, the simplest atom)

## Problems to face :

- Positronium is a neutral atom
- Positronium has a very short lifetime


Detection of the interference pattern is not going to be easy. Ionization required.

Excitation on Rydberg state is necessary.  
Laser excitation required.

## QUPLAS – I physics program:

- Quantum Decoherence with an unusual system
- Wave function interpretation
- Optical gratings and laser interaction and cooling

# Positronium fall (QUPLAS-II)



QUPLAS – I : detecting an interferometry pattern in the Talbot mode

QUPLAS – II : detecting a (gravity induced) SHIFT in the interferometry pattern

Gravity shift of 4 micron over the distance of a meter (for  $10^3$  m/s positronium)



# Measuring gravity



Methods to measure the Interference Pattern Shift

1. Quantum Talbot Interferometry

2. Moiré classical deflectometry

$$v = 800 \text{ m/s} \quad \lambda_{\text{Ps}} = 454 \text{ nm}$$


$$L^{(\text{TOT})} \approx 1 \text{ m} \quad a = 9.81 \text{ m/s}^2$$

	Talbot	Moiré
<p><b>Symmetric</b></p> 	$L^{(\text{TOT})} = 2 \frac{d_2^2}{\lambda} = 2T_L$ $d_1 = d_2 = d_3 = 476 \mu\text{m}$ $\tau = 6.25 \cdot 10^{-4} \text{ s}$ $\Delta x = 3.83 \mu\text{m}$ $\Delta x/d_3 = 8.0 \cdot 10^{-3}$	$d_1 = d_2 = d_3 \gtrsim 800 \mu\text{m}$ $L^{(\text{TOT})} = 1 \text{ m} \gg 2T_L$ $\tau = 6.25 \cdot 10^{-4} \text{ s}$ $\Delta x = 3.83 \mu\text{m}$ $\Delta x/d_3 \lesssim 4.0 \cdot 10^{-3}$
<p><b>Asymmetric</b> (<math>\eta = 3</math>)</p> 	$L^{(\text{TOT})} = \frac{(\eta + 1)^2}{\eta} T_L$ $d_2 = 291 \mu\text{m}$ $d_3 = \eta d_1 = \eta d_2 \frac{(\eta + 1)}{\eta}$ $\tau = \frac{d_1 T_L}{d_2 v} = 3.1 \cdot 10^{-4} \text{ s}$ $\Delta x = a \frac{\eta(\eta + 1)}{2} \tau^2 = 5.70 \mu\text{m}$ $\Delta x/d_3 = 4.8 \cdot 10^{-3}$	$d_2 \gtrsim 800 \mu\text{m}$ $d_3 = \eta d_1 = \eta d_2 \frac{(\eta + 1)}{\eta}$ $L^{(\text{TOT})} = 1 \text{ m} \rightarrow L = 0.25 \text{ m}$ $\tau = \frac{L}{v} = 3.1 \cdot 10^{-4} \text{ s}$ $\Delta x = a \frac{\eta(\eta + 1)}{2} \tau^2 = 5.70 \mu\text{m}$ $\Delta x/d_3 \lesssim 2 \cdot 10^{-3}$

- S. Sala et al, J. of Phys. B 48 (2015) 195002
- S. Sala et al, Phys. Rev. A in press (arxiv:1601.06539)

# Moiré deflectometry

Tested with antiprotons (in an inhomogeneous magnetic field) in the AEGLS experiment at CERN. Grating system followed by the emulsion detector.



Reference pattern obtained by light: Talbot-Lau interferometry (rephasing of light pattern at discrete distances after the grating)


$$\Delta y = \frac{F_y}{m} t^2$$

$$L_T = \frac{2d^2}{\lambda}$$

$$\Delta y_{mean} = 10.0 \pm 0.9 \pm 6.3\ \mu\text{m}$$

$$v = 4.5 \times 10^6\ \text{m/s}$$

$$t = 5.6\ \text{ns}$$



S. Aghion et al, Nature Comm. 5 (2014) 453

# Gravity measurement count rate

Count rate for a typical gravity experiment

50 mCi source with a RGM moderator (0.4% efficiency)  $\rightarrow 7 \times 10^6$  e+/s

e+  $\rightarrow$  Ps conversion (10%) and reemission (30%) by converters  $\rightarrow 2 \times 10^5$ /s

Ps solid angle of emission and interferometer geometry (0.1%)  $\rightarrow 200$ /s

Ps excitation efficiency is high but the spectral selection will introduce 10%  $\rightarrow 20$ /s

Transparency of the gratings 25%  $\rightarrow 5$ /s

## Sensitivity to g for Ps (only Talbot, interferometric methods)

Given 0.5 dots/s on the emulsion, one has, for a very realistic 50% contrast,  $d_3 = 476 \mu\text{m}$ ,  $\Delta x = 4 \mu\text{m}$  :

$$\frac{\sigma(g)}{g} \cong \frac{1}{2\pi C} \frac{1}{\sqrt{N}} \frac{d_3}{\Delta x(g)}$$

(only statistical error)

- 2% in a WEEK
- 1% in a MONTH
- <0.1% in a YEAR

# Conclusions

- Quantum Interference to explore new physics with  $e^+/e^-/Ps$
- QUPLAS is a staged project to tackle these subjects
- QUPLAS-0 : Positron and Electron quantum interference (CPT test).
- QUPLAS-I : Positronium Quantum Interferometry.
- QUPLAS-II : Positronium Gravity as a test of the Weak Equivalence Principle.

- S. Sala et al, J. of Phys. B 48 (2015) 195002
- S. Aghion et al, Nature Comm. 5 (2014) 4538
- S. Aghion et al, J. of Instr. 11 (2016) P06017
- S. Sala et al, Phys. Rev. A in press (arxiv:1601.06539)

Thank you for your attention !

## Backup Slides

# The QUPLAS Laser system for velocity-selective excitation of a continuous Ps beam

Excitation of Rydberg states :

$1 \rightarrow 2$  243 nm       $2 \rightarrow n$  ~ 732 nm


$$\tau \approx n^3 l^2$$

$$v = 10^3 \text{ m/s}$$


$$L(\text{tot}) = 1 \text{ m}$$

$$\tau \sim 1 \text{ ms}$$


$$n \sim 80-90$$



# The QUPLAS system for velocity-selective excitation of a continuous Ps beam



# The third method to measure gravity («space – gamma»)



3. Simple method of tracing a given velocity selected by the laser (highly efficient on NaI)

Three different methods to measure g	
1. Talbot interferometry	Different pitches for the gratings but similar systematics and blank measurements
2. Moiré deflectometry	
3. Space-gamma	Entirely different systematics





9/6/2016

FOMO - September 2016

33